

Patient: [REDACTED]

Reference: [REDACTED]

Patient Details

First name: [REDACTED] NHI number: [REDACTED]
Last name: [REDACTED] Date of birth: [REDACTED]
Gender: Male Telephone: [REDACTED]

Patient Address: [REDACTED]
ACC45/Claim Number: [REDACTED]

Referral Details

HealthLink Account: [REDACTED] Service Provider: [REDACTED]
Message Date: 22/2/2021 08:00:55 Date Collected: 21/2/2021 14:50

Report

Observation Reference: [REDACTED]
Observation Description: XE53L Ankle, Left (three or more vie
Relevant DateTime: 21/2/2021 14:50
Interpreted By: [REDACTED]

Result

Observation Result ID: REPORT
Observation Result Description: REPORT
Observation Result Value: ANKLE, LEFT (THREE OR MORE VIEWS)

INDICATION

Wrenching injury, bilateral malleolus pain, the lateral more than the medial.

FINDINGS

A posterolateral bone fragment over the lateral malleolus is present with bone lucency consistent with donor site. Features consistent with cortical avulsion fracture. Overlying soft tissue swelling is present. An irregular bone body with no cortical margin is projected between the medial malleolus and medial articular margin of the talus. There is cortical irregularity of the anterior articular margin of the talus. Features are consistent with second cortical avulsion fracture, probably from the talus. Soft tissue swelling is present over the medial malleolus. The distal tibia and distal fibula are otherwise intact, the distal tibiofibular syndesmosis is intact. Ankle joint is in normal alignment.

CONCLUSION

Cortical avulsion fractures of the lateral malleolus and medial aspect of the talus.